

GAITHERSBURG, Md., Sept. 2, 2022 [/PRNewswire/](#) -- Novavax, Inc. (Nasdaq: NVAX), a biotechnology company dedicated to developing and commercializing next-generation vaccines for serious infectious diseases, today announced that Swissmedic, the Swiss Agency for Therapeutic Products, has expanded its temporary authorization of Nuvaxovid™ (NVX-CoV2373) COVID-19 vaccine in Switzerland for active immunization to prevent coronavirus disease 2019 (COVID-19) caused by the severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) in adolescents aged 12 through 17 and as a heterologous and homologous booster dose for adults aged 18 and older.

"We are pleased to offer the first protein-based COVID-19 vaccine for use both in adolescents and as a booster in adults in Switzerland," said Stanley C. Erck, President and Chief Executive Officer, Novavax. "As we continue to explore best practices for managing COVID-19 long term, we have ongoing trials further exploring Nuvaxovid's efficacy and safety as a booster and preclinical data suggest that our vaccine induces immune response against Omicron variants, including BA.4/5."

Adolescents Aged 12 Through 17

The authorization for adolescents aged 12 through 17 is based on data from the ongoing [pediatric expansion](#) of the Phase 3 PREVENT-19 trial of 2,247 adolescents aged 12 through 17 years across 73 sites in the U.S. to evaluate the safety, effectiveness (immunogenicity), and efficacy of Nuvaxovid. In the pediatric expansion, Nuvaxovid achieved its primary efficacy endpoint and demonstrated 80% clinical efficacy overall at a time when the Delta variant was the predominant circulating SARS-CoV-2 strain in the U.S.

Preliminary safety data from the pediatric expansion showed the vaccine to be generally well-tolerated. Serious and severe adverse events were low in number and balanced between vaccine and placebo groups, and not considered related to the vaccine. Local and systemic reactogenicity was generally lower than or similar to adults, after the first and second dose. The most common adverse reactions observed were injection site tenderness/pain, headache, myalgia, fatigue, and malaise. There was no increase in reactogenicity in younger (12 to <15 years old) adolescents compared to older (15 to <18 years old) adolescents. No new safety signal was observed through the placebo-controlled portion of the pediatric expansion.

Booster in Adults Aged 18 and Older

The authorization for the booster dose in adults aged 18 and older is supported by data from Novavax' Phase 2 trial conducted in Australia, from a separate [Phase 2 trial](#) conducted in South Africa, and from the UK-sponsored COV-BOOST trial. As part of the Phase 2 trials, a single booster dose of Nuvaxovid was administered to healthy adult participants approximately six months after their primary two-dose vaccination series of Nuvaxovid. The third dose produced increased immune responses comparable to or exceeding levels associated with protection in Phase 3 clinical trials. In the COV-BOOST trial, Nuvaxovid induced a meaningful antibody response when used as a heterologous third booster dose.

In the Novavax-sponsored trials, following the booster, local and systemic reactions had a median duration of approximately two days. The incidence of Grade 3 or higher events remained relatively low. Safety reporting of reactogenicity events showed an increasing incidence across all three doses of Nuvaxovid, often seen with increased immunogenicity. Medically attended adverse events, potentially immune-mediated medical conditions, and severe adverse events occurred infrequently following the booster dose and were balanced between vaccine and placebo groups.

In the 12 through 17 year-old population, NVX-CoV2373 has also been granted authorization in the [U.S.](#), the [European Union](#), the [United Kingdom](#), [Australia](#), [New Zealand](#), [Japan](#), [Thailand](#), [India](#), and [South Korea](#). The vaccine has also been authorized in [Japan](#), [Australia](#), and [New Zealand](#) as a booster. Nuvaxovid is actively under review for both indications in other markets.

Swissmedic granted temporary authorization in [April 2022](#) for use of Nuvaxovid in adults aged 18 and older.

Trade Name in the U.S.

The trade name Nuvaxovid™ has not yet been approved by the U.S. Food and Drug Administration.

Authorized Use

Nuvaxovid is indicated for active immunisation to prevent COVID-19 caused by SARS-CoV-2 in individuals 12 years of age and older. The use of this vaccine should be in accordance with official recommendations.

Important Safety Information: Switzerland

- Nuvaxovid is contraindicated in persons who have a hypersensitivity to the active substance, or to any of the excipients.
- Events of anaphylaxis have been reported with administration of COVID-19 vaccines including Nuvaxovid.

Appropriate medical treatment and supervision should be available in case of an anaphylactic reaction following the administration of the vaccine. Close observation for at least 15 minutes is recommended and a second dose of the vaccine should not be given to those who have experienced anaphylaxis to the first dose of Nuvaxovid.

- Very rare cases of myocarditis and pericarditis have been reported following the use of Nuvaxovid. Healthcare professionals should be alert to the signs and symptoms of myocarditis and pericarditis. Vaccinees (including parents or caregivers) should be instructed to seek immediate medical attention if they develop symptoms indicative of myocarditis or pericarditis such as (acute and persisting) chest pain, shortness of breath, or palpitations following vaccination. Healthcare professionals should consult guidance and/or specialists to diagnose and treat this condition.
- Anxiety-related reactions, including vasovagal reactions (syncope), hyperventilation, or stress-related reactions may occur in association with vaccination as a psychogenic response to the needle injection. It is important that precautions are in place to avoid injury from fainting.
- Vaccination should be postponed in individuals suffering from an acute severe febrile illness or acute infection. The presence of a minor infection and/or low-grade fever should not delay vaccination.
- Nuvaxovid should be given with caution in individuals receiving anticoagulant therapy or those with thrombocytopenia or any coagulation disorder (such as haemophilia) because bleeding or bruising may occur following an intramuscular administration in these individuals.
- The efficacy of Nuvaxovid may be lower in immunosuppressed individuals.
- Administration of Nuvaxovid in pregnancy should only be considered when the potential benefits outweigh any potential risks for the mother and foetus.
- The effects with Nuvaxovid may temporarily affect the ability to drive or use machines.
- The duration of protection afforded by the vaccine is unknown as it is still being determined by ongoing clinical trials. Individuals may not be fully protected until 7 days after their second dose. As with all vaccines, vaccination with Nuvaxovid may not protect all vaccine recipients.
- Very common ($\geq 1/10$) and common ($\geq 1/100$ to $< 1/10$) adverse reactions observed during clinical studies in individuals 12 years of age and older were headache, nausea or vomiting, myalgia, arthralgia, injection site tenderness/redness, pyrexia. Fever was observed more frequently in adolescents aged 12 through 17 years compared to adults, with the frequency being very common after the second dose in adolescents.

For more information on Nuvaxovid, including the Summary of Product Characteristics with Package Leaflet, adverse event reporting instructions, or to request additional information, please visit the following website:

- [Swissmedic](#)

About Nuvaxovid™ (NVX-CoV2373)

Nuvaxovid is a protein-based vaccine engineered from the genetic sequence of the first strain of SARS-CoV-2, the virus that causes COVID-19 disease. The vaccine was created using Novavax' recombinant nanoparticle technology to generate antigen derived from the coronavirus S protein and is formulated with Novavax' patented saponin-based Matrix-M™ adjuvant to enhance the immune response and stimulate high levels of neutralizing antibodies. Nuvaxovid contains purified protein antigen and can neither replicate, nor can it cause COVID-19.

Nuvaxovid is packaged as a ready-to-use liquid formulation in a vial containing ten doses. The vaccination regimen calls for two 0.5 ml doses (5 mcg antigen and 50 mcg Matrix-M adjuvant) given intramuscularly 21 days apart. The vaccine is stored at 2°- 8° Celsius, enabling the use of existing vaccine supply and cold chain channels. Use of the vaccine should be in accordance with official recommendations.

Novavax has established partnerships for the manufacture, commercialization, and distribution of Nuvaxovid worldwide. Existing authorizations leverage Novavax' manufacturing partnership with Serum Institute of India, the world's largest vaccine manufacturer by volume. They are being supplemented with data from additional manufacturing sites throughout Novavax' global supply chain.

About the NVX-CoV2373 Phase 3 Trials

NVX-CoV2373 continues being evaluated in two pivotal Phase 3 trials. PREVENT-19 (the **PRE**-fusion protein subunit **V**accine **E**fficacy **N**ovavax **T**rial | COVID-19) is a 2:1 randomized, placebo-controlled, observer-blinded trial to evaluate the efficacy, safety, and immunogenicity of NVX-CoV2373 with Matrix-M adjuvant in 29,960 participants 18 years of age and over in 119 locations in the U.S. and Mexico. The primary endpoint for PREVENT-19 was the first occurrence of PCR-confirmed symptomatic (mild, moderate, or severe) COVID-19 with onset at least seven days after the second dose in serologically negative (to SARS-CoV-2) adult participants at baseline. The statistical success criterion included a lower bound of 95% CI $> 30\%$. A secondary endpoint was the prevention of PCR-confirmed, symptomatic moderate or severe COVID-19. Both endpoints were assessed at least seven days after the second study vaccination in volunteers who had not been previously infected with SARS-CoV-2. In the trial, NVX-CoV2373 achieved 90.4% efficacy overall. It was generally well-tolerated and elicited a robust antibody response after the second dose in both studies. Full results of the trial were published

in the [*New England Journal of Medicine*](#) (*NEJM*).

The pediatric expansion of PREVENT-19 is a 2:1 randomized, placebo-controlled, observer-blinded trial to evaluate the safety, effectiveness, and efficacy of NVX-CoV2373 with Matrix-M adjuvant in 2,247 adolescent participants 12 to 17 years of age in 73 locations in the U.S., compared with placebo. In the pediatric trial, NVX-CoV2373 achieved its primary effectiveness endpoint (non-inferiority of the neutralizing antibody response compared to young adult participants 18 through 25 years of age from PREVENT-19) and demonstrated 80% efficacy overall at a time when the Delta variant of concern was the predominant circulating strain in the U.S. Additionally, immune responses were about two-to-three-fold higher in adolescents than in adults against all variants studied.

Additionally, a trial conducted in the U.K. with 14,039 participants aged 18 years and over was designed as a randomized, placebo-controlled, observer-blinded study and achieved overall efficacy of 89.7%. The primary endpoint was based on the first occurrence of PCR-confirmed symptomatic (mild, moderate, or severe) COVID-19 with onset at least seven days after the second study vaccination in serologically negative (to SARS-CoV-2) adult participants at baseline. Full results of the trial were published in [*NEJM*](#).

About Matrix-M™ Adjuvant

Novavax' patented saponin-based Matrix-M adjuvant has demonstrated a potent and well-tolerated effect by stimulating the entry of antigen-presenting cells into the injection site and enhancing antigen presentation in local lymph nodes, boosting immune response.

About Novavax

Novavax, Inc. (Nasdaq: NVAX) is a biotechnology company that promotes improved health globally through the discovery, development, and commercialization of innovative vaccines to prevent serious infectious diseases. The company's proprietary recombinant technology platform harnesses the power and speed of genetic engineering to efficiently produce highly immunogenic nanoparticles designed to address urgent global health needs. NVX-CoV2373, the company's COVID-19 vaccine, has received conditional authorization from multiple regulatory authorities globally, including the U.S., European Commission, and the World Health Organization. The vaccine is currently under review by multiple regulatory agencies worldwide and is under review in the U.S. for use in adolescents and as a booster. In addition to its COVID-19 vaccine, Novavax is also currently evaluating a COVID-seasonal influenza combination vaccine candidate in a Phase 1/2 clinical trial, which combines NVX-CoV2373 and NanoFlu*, its quadrivalent influenza investigational vaccine candidate, and is also evaluating an Omicron strain-based vaccine (NVX-CoV2515) as well as a bivalent format Omicron-based / original strain-based vaccine. These vaccine candidates incorporate Novavax' proprietary saponin-based Matrix-M adjuvant to enhance the immune response and stimulate high levels of neutralizing antibodies.

For more information, visit www.novavax.com and connect with us on [LinkedIn](#).

*NanoFlu identifies a recombinant hemagglutinin (HA) protein nanoparticle influenza vaccine candidate produced by Novavax. This investigational candidate was evaluated during a controlled phase 3 trial conducted during the 2019-2020 influenza season.

Forward-Looking Statements

Statements herein relating to the future of Novavax, its operating plans and prospects, its partnerships, the timing of clinical trial results, the ongoing development of NVX-CoV2373, a COVID-seasonal influenza investigational vaccine candidate, the scope, timing and outcome of future regulatory filings and actions, including with respect to an FDA EUA decision and potential CDC recommendation for NVX-CoV2373, Novavax' plans to supplement existing authorizations with data from the additional manufacturing sites in Novavax' global supply chain, additional worldwide authorizations of NVX-CoV2373 for use in adults and adolescents and as a booster, the potential impact and reach of Novavax and NVX-CoV2373 in addressing vaccine access, controlling the pandemic and protecting populations, and the efficacy, safety and intended utilization of NVX-CoV2373 and expected administration of NVX-CoV2373 are forward-looking statements. Novavax cautions that these forward-looking statements are subject to numerous risks and uncertainties that could cause actual results to differ materially from those expressed or implied by such statements. These risks and uncertainties include, without limitation, challenges satisfying, alone or together with partners, various safety, efficacy, and product characterization requirements, including those related to process qualification and assay validation, necessary to satisfy applicable regulatory authorities; difficulty obtaining scarce raw materials and supplies; resource constraints, including human capital and manufacturing capacity, on the ability of Novavax to pursue planned regulatory pathways; challenges meeting contractual requirements under agreements with multiple commercial, governmental, and other entities; and those other risk factors identified in the "Risk Factors" and "Management's Discussion and Analysis of Financial Condition and Results of Operations" sections of Novavax' Annual Report on Form 10-K for the year ended December 31, 2021 and subsequent Quarterly Reports on Form 10-Q, as filed with the Securities and Exchange Commission (SEC). We caution investors not to place considerable reliance on forward-looking statements contained in this press release. You are encouraged to read

our filings with the SEC, available at www.sec.gov and www.novavax.com, for a discussion of these and other risks and uncertainties. The forward-looking statements in this press release speak only as of the date of this document, and we undertake no obligation to update or revise any of the statements. Our business is subject to substantial risks and uncertainties, including those referenced above. Investors, potential investors, and others should give careful consideration to these risks and uncertainties.

Contacts:

Investors

Erika Schultz | 240-268-2022

ir@novavax.com

Media

Ali Chartan or Giovanna Chandler | 202-709-5563

media@novavax.com

SOURCE Novavax, Inc.

<https://ir.novavax.com/2022-09-02-Swissmedic-Authorizes-Novavax-Nuvaxovid-TM-COVID-19-Vaccine-for-Adolescents-Aged-12-Through-17-and-as-a-Booster-in-Adults-Aged-18-and-Older>